

Conference Abstract


http://dx.doi.org/10.21065/AdvFoodNutrSci.4.19

9th Pak Pharma Healthcare Expo

1ST Pakistan Food –Nutrition International Summit 2019

Monday –Thursday, July 10-11, 2019

Pak-China Friendship Centre, Garden Avenue, Islamabad, Pakistan



Conference Abstract

HYPOCHOLESTEROLEMIC EFFECT OF DESIGNER YOGURTS FORTIFIED WITH OMEGA FATTY ACIDS AND DIETARY FIBRES IN HYPERCHOLESTEROLEMIC SUBJECTS

Nazir Ahmad¹, Hafsa Mughal², Muhammad Faisal Manzoor⁴, Tariq Ismail³, Mahr un Nisa¹

1. Institute of Home and Food Sciences, Faculty of Life Sciences, Government College University, Faisalabad-Pakistan
2. Department Nutrition, Aziz Fatima Hospital, Faisalabad-Pakistan
3. Institute of Food Science and Nutrition, Bahauddin Zakariya University, Multan, Pakistan
4. School of Food Science and Engineering, South China University of Technology, Guangzhou-China

Corresponding author:

Nazir Ahmad. Institute of Home and Food Sciences, Faculty of Life Sciences, Government College University, Faisalabad-Pakistan
P: +92 333 06 12 809
E: drnazirahmad@gcuf.edu.pk

Citation:

Nazir A, Hafsa M, Muhammad Faisal M, Tariq I, Mahr un Nisa. Hypcholesterolemic Effect of Designer Yogurts Fortified with Omega Fatty Acids and Dietary Fibres in Hypercholesterolemic Subjects. Adv Food Nutr Sci. Vol. 4. 2019. p 19-20

Funding:

The authors received no direct funding for this research.

Competing Interests:

The authors declare no competing interests

Additional information is available at the end of the article.

Abstract

Background: The saturated fatty acids of dairy products have negative impact on plasma cholesterol and limited the consumption and benefits of these products for human. The aim of this study was to evaluate the effect of designer yogurts with improved health lipid indexes through double fortification with omega fatty acids and fibres of extruded flaxseed powder (EFSP) in hypercholesterolemic subjects.

Methods: A cohort study of 90 days was designed to explore the effect of yogurts on hypercholesterolemia. The hypercholesterolemic subjects were randomly divided in group 1 and group 2 and blood samples and blood pressure were taken at 0 day (baseline). Group 1 was offered plain sheep yogurt and group 2 cow milk yogurt for 30 days (intervention). At the end 30th day, blood samples and blood pressure were taken. Participants were allowed to eat routine diet for further 30 days (wash-out period). After wash-out periods, the blood samples and blood pressure were taken (baseline) and EFSP fortified sheep and cow milk yogurts were offered to group 1 and 2, respectively, for next 30 days (intervention). At the end, blood samples and blood pressure were recorded. Descriptive statistics was used to compare the influence of yogurts intake on hyperlipidemia

Results: Significant increase in serum TC (6.33 ± 0.48 to 6.51 ± 0.04) and LDL-C (5.17 ± 0.28 to 5.37 ± 0.47) was observed in participants consumed plain sheep milk yoghurt for 30 days and non-significant change was observed for HDL-C and TG. After washout period of 30

days, no significant change was observed in any cholesterol or TG. Significant decrease in TC (6.47 ± 0.95 to 6.28 ± 0.84) and LDL-C (5.35 ± 0.29 to 5.13 ± 0.44) was observed while TG and HDL-C did not change significantly in group 1 consuming EFSP fortified yogurt for 30 days. Non-significant change was observed for serum TC, HDL-C, LDL-C and TG in group 2 consuming plain cow milk yogurt. After washout period, significant decrease in serum TC (6.38 ± 1.01 to 6.20 ± 0.98), HDL-C (1.10 ± 0.21 to 1.27 ± 0.46) and LDL-C (5.30 ± 1.16 to 4.98 ± 0.99) was noticed while TG level did not change significantly in group 2 consuming EFSP fortified cow milk yogurt. The intake of fortified yogurts reduced pressure significantly blood pressure in both groups.

Conclusions: The results show that regular dietary intake of yogurt double fortified with omega fatty acids and fibres can help to reduce the cholesterol and elevated blood pressure in hypercholesterolemic subjects

Key words: Milk yogurt, omega fatty acids, dietary fibres, flaxseed, fortification, hypercholesterolemia



© 2019 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits

The abstract appearing is approved to present in 9th Pak Pharma Healthcare Expo | 1ST Pakistan Food –Nutrition International Summit 2019, November 25, 2018 at 1194 Matheson Blvd E Mississauga, ON L4W 1Y2 Canada. The submissions were selected and processed by the Organizing Committee to be presented in oral or poster session, and were subject to review by the Program Committee.